

Linearized solutions of the Einstein equations within a Bondi-Sachs framework, and implications for boundary conditions in numerical simulations

Nigel T. Bishop

Department of Mathematical Sciences, University of South Africa, P.O. Box 392, Unisa 0003, South Africa
(Dated: 14 March 2005)

We linearize the Einstein equations when the metric is Bondi-Sachs, when the background is Schwarzschild or Minkowski, and when there is a matter source in the form of a thin shell whose density varies with time and angular position. By performing an eigenfunction decomposition, we reduce the problem to a system of linear ordinary differential equations which we are able to solve. The solutions are relevant to the characteristic formulation of numerical relativity: (a) as exact solutions against which computations of gravitational radiation can be compared; and (b) in formulating boundary conditions on the $r = 2M$ Schwarzschild horizon.

PACS numbers: 04.25.Nx, 04.25.Dm, 04.30.Db

I. INTRODUCTION

This work is motivated by the needs of numerical relativity, specifically the characteristic, or null cone, formulation used in the PITT code [1, 2] (and see also [3, 4, 5, 6]). It should be noted that the PITT code is not the only approach to characteristic numerical relativity – for example, see [7, 8, 9, 10, 11, 12]. The PITT code exhibits long-term stability in evolving single black hole spacetimes [13], and has also been successfully applied to the problem of nonlinear scattering of gravitational radiation by a black hole [14]. However, it has not been possible to compute gravitational radiation emitted in astrophysically interesting scenarios involving matter – for example, a star in close orbit around a black hole [5]. Variations of the method of calculating the gravitational radiation have been investigated [15] and [14, 16], but the problem has not been resolved. An important difficulty in trying to analyze the problem is that, while methods such as the quadrupole formula indicate approximately what the final result of the computation should be, there is no guidance concerning the intermediate quantities. Thus, in this paper we construct analytic solutions containing gravitational radiation and matter, against which a numerical code can be tested.

In addition, we discover an unexpected issue concerning the boundary condition commonly imposed on a black hole horizon – the details are given in Sec. V A.

Linearization and eigenfunction decomposition are tools commonly used to obtain an analytic understanding of mathematically complex systems. The Einstein equations are a particularly complicated system of nonlinear partial differential equations, and there is an extensive literature on linearization techniques. Work on linearization within the characteristic formalism includes the Newtonian correspondence method [17, 18, 19], which is related to linearization about a Minkowski background. More recently [20, 21], the Teukolsky equation [22] has been used as a starting point to investigate gravitational radiation on a Schwarzschild background; but we do not use that formalism here because the coordinates are not those of the PITT code. The key point about the approach presented here is that we use the same coordinates and metric variables as in the code, so that comparisons between numerical and analytic solutions can easily be made. We also allow for the presence of a matter source.

We seek a physical situation that is without symmetry and that radiates gravitationally, but in which the linearized problem has a solution in simple analytic form. The model problem should be relevant to the physical problem of a star in quasi-circular orbit around a black hole. We achieve these goals by considering a Schwarzschild black hole of mass M (with the case $M = 0$ corresponding to a Minkowski background), surrounded by a thin spherical shell of matter whose density varies in time u as $e^{i\nu u}$ (with the static case recovered by setting $\nu = 0$), and whose angular dependence is that of a spherical harmonic. We find a solution for all metric variables in simple analytic form, except when $M \neq 0$ and $\nu \neq 0$ in which case solutions are obtained as power series.

The calculations reported in this paper are very long, and would have been impossible to perform without the use of a computer algebra system, in this case Maple.

In Sec. II we give a summary of the formalism concerning the Bondi-Sachs metric in numerical relativity, together with the formalism of spin-weighted spherical harmonics. Next, in Sec. III, we specify the model problem to be solved and work out the resulting Einstein equations, reducing the problem to coupled linear ordinary differential equations. Solutions are found in Sec. IV, and presented according to the values of the parameters ν and M . The paper concludes, in Sec. V, with a discussion of appropriate boundary conditions at a black hole horizon, and an outline of opportunities for taking this work further.

II. FORMALISM

A. Bondi-Sachs metric

The formalism for the numerical evolution of Einstein's equations, in null cone coordinates, is well known [1, 2, 3, 4, 23, 24]. For the sake of completeness, we give a summary of those aspects of the formalism that will be used here. We start with coordinates based upon a family of outgoing null hypersurfaces. We let u label these hypersurfaces, x^A ($A = 2, 3$), label the null rays and r be a surface area coordinate. In the resulting $x^\alpha = (u, r, x^A)$ coordinates, the metric takes the Bondi-Sachs form [24, 25]

$$ds^2 = - \left(e^{2\beta} \left(1 + \frac{W}{r} \right) - r^2 h_{AB} U^A U^B \right) du^2 - 2e^{2\beta} dudr - 2r^2 h_{AB} U^B dudx^A + r^2 h_{AB} dx^A dx^B, \quad (1)$$

where $h^{AB} h_{BC} = \delta_C^A$ and $\det(h_{AB}) = \det(q_{AB})$, with q_{AB} a unit sphere metric. We work in stereographic coordinates $x^A = (q, p)$ for which the unit sphere metric is

$$q_{AB} dx^A dx^B = \frac{4}{P^2} (dq^2 + dp^2), \quad (2)$$

where

$$P = 1 + q^2 + p^2. \quad (3)$$

We also introduce a complex dyad q_A defined by

$$q^A = \frac{P}{2} (1, i), \quad q_A = \frac{2}{P} (1, i) \quad (4)$$

with $i = \sqrt{-1}$. For an arbitrary Bondi-Sachs metric, h_{AB} can then be represented by its dyad component

$$J = h_{AB} q^A q^B / 2, \quad (5)$$

with the spherically symmetric case characterized by $J = 0$. We introduce the spin-weighted field

$$U = U^A q_A, \quad (6)$$

as well as the (complex differential) eth operators $\bar{\eth}$ and $\eth$ (see [26] for full details).

Einstein's equations $R_{\alpha\beta} = 8\pi(T_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}T)$ are classified as: hypersurface equations $-R_{11}, q^A R_{1A}, h^{AB} R_{AB}$ – forming a hierarchical set for β, U and W ; evolution equation $q^A q^B R_{AB}$ for J ; and constraints $R_{0\alpha}$. An evolution problem is normally formulated in the region of spacetime between a timelike or null worldtube Γ and future null infinity, with (free) initial data J given on $u = 0$. Boundary data for β, U, W, J that satisfies the constraints, is required on Γ . In the case that the spacetime contains a single Schwarzschild black hole, Γ is usually taken to be the (past) event horizon $r = 2M$.

In a general situation the various metric components do not admit simple physical interpretations, but nevertheless a review of some particular cases does illustrate the roles of the various coefficients. Minkowski spacetime is recovered by setting all the coefficients to zero, i.e. $\beta = W = U = J = 0$. Schwarzschild spacetime has $J = U = 0$, and is usually described with $\beta = 0$, $W = -2M$, but can also be represented by $\beta = \beta_c$ (constant) and $W = (e^{2\beta_c} - 1)r - 2M$. The quantities J and U are zero when there is spherical symmetry, and so can be regarded as a measure of the deviation from spherical symmetry: the quantities are interlinked, and in the linearized regime contain all the dynamic content of the gravitational field.

B. Spin-weighted spherical harmonics

We will be using spin-weighted spherical harmonics [27, 28] using the formalism described in [14]. It will prove convenient to use ${}_s Z_{\ell m}$ rather than the usual ${}_s Y_{\ell m}$ as basis functions, where

$$\begin{aligned} {}_s Z_{\ell m} &= \frac{1}{\sqrt{2}} [{}_s Y_{\ell m} + (-1)^m {}_s Y_{\ell -m}] \text{ for } m > 0 \\ {}_s Z_{\ell m} &= \frac{i}{\sqrt{2}} [(-1)^m {}_s Y_{\ell m} - {}_s Y_{\ell -m}] \text{ for } m < 0 \\ {}_s Z_{\ell 0} &= {}_s Y_{\ell 0}, \end{aligned} \quad (7)$$

and note that [14] uses the notation ${}_s R_{\ell m}$ rather than the ${}_s Z_{\ell m}$ used here; we use a different notation to avoid any confusion with the Ricci tensor. In the case $s = 0$, the s will be omitted, i.e. $Z_{\ell m} = {}_0 Z_{\ell m}$. Some properties of the ${}_s Z_{\ell m}$ that will be used later are

- They are orthonormal
- They are **real**, which is the crucial point – quantities like $\bar{\partial}U + \bar{\partial}\bar{U}$ do not introduce m and $-m$ mode mixing.
- Defining

$$L_2 = -\ell(\ell + 1) \quad (8)$$

then

$$\bar{\partial}\bar{\partial}Z_{\ell m} = \bar{\partial}\bar{\partial}Z_{\ell m} = L_2 Z_{\ell m}, \quad \bar{\partial}\bar{\partial}^2 Z_{\ell m} = (L_2 + 2)\bar{\partial}Z_{\ell m}, \quad \bar{\partial}^2\bar{\partial}^2 Z_{\ell m} = \bar{\partial}^2\bar{\partial}^2 Z_{\ell m} = L_2(L_2 + 2)Z_{\ell m}. \quad (9)$$

- The effect of the $\bar{\partial}$ operator acting on $Z_{\ell m}$ is

$$\bar{\partial}Z_{\ell m} = \sqrt{-L_2} {}_{-1}Z_{\ell m}, \quad \bar{\partial}^2 Z_{\ell m} = \sqrt{-(\ell - 1)L_2(\ell + 2)} {}_{-2}Z_{\ell m}, \quad (10)$$

III. SIMPLIFICATION OF THE EINSTEIN EQUATIONS FOR A BONDI-SACHS METRIC

A. The linearized Einstein equations

Physically, we consider a matter distribution in the form of a thin, low-density, shell around either a Schwarzschild black hole or empty space. The shell is spherical at $r = \text{constant}$, but the density ρ can vary around the shell so the problem is not spherically symmetric. We regard the density and metric quantities as being small, i.e.

$$\rho, J, \beta, U, w = \mathcal{O}(\epsilon) \text{ where } W = -2M + w, \quad (11)$$

with M the mass of the Schwarzschild black hole (and, of course, the Minkowski case is recovered simply by setting $M = 0$). All terms in the Einstein equations of order $\mathcal{O}(\epsilon^2)$ will be set to zero. That is, we perform a standard linearization of the Einstein equations.

Using the above ansatz, we found the following forms for the hypersurface equations and the evolution equation

$$R_{11} : \quad \frac{4}{r}\beta_{,r} = 8\pi T_{11} \quad (12)$$

$$q^A R_{1A} : \quad \frac{1}{2r} (4\bar{\partial}\beta - 2r\bar{\partial}\beta_{,r} + r\bar{\partial}J_{,r} + r^3 U_{,rr} + 4r^2 U_{,r}) = 8\pi q^A T_{1A} \quad (13)$$

$$h^{AB} R_{AB} : \quad (4 - 2\bar{\partial}\bar{\partial})\beta + \frac{1}{2}(\bar{\partial}^2 J + \bar{\partial}^2 \bar{J}) + \frac{1}{2r^2}(r^4 \bar{\partial}\bar{U} + r^4 \bar{\partial}U)_{,r} - 2w_{,r} = 8\pi(h^{AB} T_{AB} - r^2 T) \quad (14)$$

$$q^A q^B R_{AB} : \quad -2\bar{\partial}^2 \beta + (r^2 \bar{\partial}U)_{,r} - 2(r - M)J_{,r} - \left(1 - \frac{2M}{r}\right)r^2 J_{,rr} + 2r(rJ)_{,ur} = 8\pi q^A q^B T_{AB}. \quad (15)$$

In addition, we evaluated the constraint equations; these will be needed only off the matter shell, and the formulas below are for the vacuum case

$$R_{00} : \quad \frac{1}{2r^3} \left(r(r - 2M)w_{,rr} + \bar{\partial}\bar{\partial}w + 2(r - 2M)\bar{\partial}\bar{\partial}\beta - Mr(\bar{\partial}\bar{U} + \bar{\partial}U) - 4r(r - 2M)\beta_{,u} - r^3(\bar{\partial}\bar{U} + \bar{\partial}U)_{,u} + 2rw_{,u} \right) = 0 \quad (16)$$

$$R_{01} : \quad \frac{1}{4r^2} \left(2rw_{,rr} + 4\bar{\partial}\bar{\partial}\beta - (r^2 \bar{\partial}\bar{U} + r^2 \bar{\partial}U)_{,r} \right) = 0 \quad (17)$$

$$q^A R_{0A} : \quad \frac{1}{4r^2} \left(2r\bar{\partial}w_{,r} - 2\bar{\partial}w + 2r^2(r - 2M)(4U_{,r} + rU_{,rr}) + 4r^2 U + r^2(\bar{\partial}\bar{\partial}U - \bar{\partial}^2 \bar{U}) + 2r^2 \bar{\partial}J_{,u} - 2r^4 U_{,ur} - 4r^2 \bar{\partial}\beta_{,u} \right) = 0. \quad (18)$$

B. Eigenfunction decomposition

We first suppose that the various metric quantities can be written as

$$J = J_0(r) \cos(\nu u) \bar{\partial}^2 Z_{\ell m}, \quad U = U_0(r) \cos(\nu u) \bar{\partial} Z_{\ell m}, \quad \beta = \beta_0(r) \cos(\nu u) Z_{\ell m}, \quad w = w_0(r) \cos(\nu u) Z_{\ell m}. \quad (19)$$

Of course, a general spacetime will not satisfy Eq. (19), but it can be represented by summing over ℓ and m and integrating over r_0 and ν . However, for our purposes we regard ℓ , m , r_0 and ν as fixed. Later, we will express the time variation in the more usual way as $e^{i\nu u}$, but for the time being we need the dependence to be explicitly real. The reason is that we are now using imaginary quantities as a representation on S^2 , and this is not related to the phase.

Having chosen basis functions that are eigenfunctions of the relevant operators, the result of applying ansatz Eq. (19) is that we reduce Eqs. (12) to (15) to ordinary differential equations

$$\frac{4}{r} \beta_{0,r} \cos(\nu u) Z_{\ell m} = 8\pi T_{11} \quad (20)$$

$$\frac{1}{2r} (4\beta_0 - 2r\beta_{0,r} + r^3 U_{0,rr} + 4r^2 U_{0,r} + (2 + L_2)rJ_0) \cos(\nu u) \bar{\partial} Z_{\ell m} = 8\pi q^A T_{1A} \quad (21)$$

$$\left(2(2 - L_2)\beta_0 + L_2(L_2 + 2)J_0 + \frac{1}{r^2} (r^4 L_2 U_0)_{,r} - 2w_{0,r} \right) \cos(\nu u) Z_{\ell m} = 8\pi (h^{AB} T_{AB} - r^2 T) \quad (22)$$

$$\left(-2\beta_0 + 2U_0 r + r^2 U_{0,r} - 2(r - M)J_{0,r} - r^2 \left(1 - \frac{2M}{r} \right) J_{0,rr} + 2r\Psi(J_0 + rJ_{0,r}) \right) \cos(\nu u) \bar{\partial}^2 Z_{\ell m} = 8\pi q^A q^B T_{AB}. \quad (23)$$

where Ψ stands for

$$\Psi = -\nu \frac{\sin(\nu u)}{\cos(\nu u)}. \quad (24)$$

The functions $\beta_0(r)$, $U_0(r)$, $w_0(r)$ and $J_0(r)$ can be taken as real in the above equations. Thus, we can now replace $\cos(\nu u)$ by the real part of $e^{i\nu u}$, and $\Psi \cos(\nu u)$ by the real part of $-i\nu e^{i\nu u}$. Any imaginary component introduced into any of $\beta_0(r)$, $U_0(r)$, $w_0(r)$ or $J_0(r)$ can now unambiguously be interpreted as a phase variation.

C. The news

Linearizing Eq. (B5) in [1], we find that the gravitational news is

$$N = \lim_{r \rightarrow \infty} \left(-\frac{1}{2} r^2 J_{,ru} + \frac{1}{2} \bar{\partial}^2 \omega + \bar{\partial}^2 \beta \right) \quad (25)$$

where ω satisfies (see Eqs. (11), (31) and (B1) of [1])

$$2(\omega^2 + \bar{\partial} \bar{\partial} \log \omega) = 2 + \frac{1}{2} (\bar{\partial}^2 J + \bar{\partial}^2 \bar{J}). \quad (26)$$

Now applying the same type of ansatz to ω as to the other metric variables, i.e.,

$$\omega = 1 + \omega_0 Z_{\ell m} e^{i\nu u} \quad (27)$$

where $\omega_0 = \mathcal{O}(\epsilon)$, we find that Eq. (26) simplifies to

$$(4\omega_0 + 2L_2\omega_0 - L_2(L_2 + 2)J_{0\infty}) Z_{\ell m} e^{i\nu u} = 0, \quad (28)$$

where $J_{0\infty}$ means

$$J_{0\infty} = \lim_{r \rightarrow \infty} J_0(r). \quad (29)$$

Solving Eq. (28) for ω_0 , we find

$$\omega_0 = \frac{L_2}{2} J_{0\infty} \quad (30)$$

so that Eq. (25) becomes

$$N = \left(-\frac{i\nu}{2} \lim_{r \rightarrow \infty} (r^2 J_{0,r}) + \frac{L_2}{4} J_{0\infty} + \beta_0 \right) \sqrt{-(\ell - 1)L_2(\ell + 2)} Z_{\ell m} e^{i\nu u}. \quad (31)$$

D. Solution procedure

Eqs. (20) to (23) may be solved in the following way. First, Eq. (20) shows that β_0 takes a constant value, say β_{0+} , in $r > r_0$, and a different constant value, say β_{0-} , in $r < r_0$ (although we will set $\beta_{0-} = 0$). Then Eqs. (21) and (23) constitute a coupled system of ordinary differential equations in $U_0(r)$ and $J_0(r)$, and once they have been solved Eq. (22) can be solved for $w_0(r)$. The computer algebra does solve the systems for general ℓ , but some of the expressions get very complicated and in order to simplify the presentation we now specialize to the case

$$\ell = 2 \quad (32)$$

so that $L_2 = -6$. In $r > r_0$, we remove all common factors from Eqs. (21) and (23) and they simplify to (with the form in $r < r_0$ obtained by replacing β_{0+} by zero, as well as J_{0+} by J_{0-} etc.)

$$4\beta_{0+} + r^3 U_{0+,rr} + 4r^2 U_{0+,r} - 4r J_{0+} = 0 \quad (33)$$

$$-2\beta_{0+} + 2U_{0+,r} + r^2 U_{0+,r} - 2(r - M)J_{0+,r} - r^2 \left(1 - \frac{2M}{r}\right) J_{0+,rr} + 2r i \nu (J_{0+} + r J_{0+,r}) = 0 \quad (34)$$

$$16\beta_{0+} + 24J_{0+} - \frac{6}{r^2} (r^4 U_{0+,r})_{,r} - 2w_{0+,r} = 0. \quad (35)$$

The constraints simplify to

$$R_{00} : \frac{1}{2r^3} \left((r^2 - 2Mr)w_{0+,rr} - 6w_{0+} - 12(r - 2M)\beta_{0+} + 12MrU_{0+} - 4r(r - 2M)i\nu\beta_{0+} + 12r^3 i\nu U_{0+} + 2r i \nu w_{0+} \right) = 0 \quad (36)$$

$$R_{01} : \frac{1}{2r^2} \left(r w_{0+,rr} - 12\beta_{0+} + 6r^2 U_{0+,r} + 12r U_{0+} \right) = 0 \quad (37)$$

$$q^A R_{0A} : \frac{1}{2r^2} \left(r w_{0+,r} - w_{0+} + 4r^3 U_{0+,r} + r^4 U_{0+,rr} + 2r^2 U_{0+} - 2Mr^3 U_{0+,rr} - 8Mr^2 U_{0+,r} - 4r^2 i \nu J_{0+} - r^4 i \nu U_{0+,r} - 2r^2 i \nu \beta_{0+} \right) = 0. \quad (38)$$

IV. SOLUTIONS

Eqs. (33) and (34) can be manipulated to give

$$-2J_2(2x + 8Mx^2 + i\nu) + 2\frac{dJ_2}{dx} (2x^2 + i\nu x - 7x^3 M) + x^3(1 - 2xM)\frac{d^2 J_2}{dx^2} = 0 \quad (39)$$

where $J_2(x) \equiv d^2 J_{0+}/dx^2$ and $x = 1/r$. Clearly, two solutions for $J_{0+}(r)$ are of the form $C_1 + C_2/r$, and the remaining two solutions depend on the values of ν and M . Once a general solution of Eq. (39) has been found, it is straightforward to calculate $U_{0+}(r)$ and $w_{0+}(r)$. We will present the solutions in terms of the four cases $\nu =, \neq 0$; $M =, \neq 0$.

A. Case $M = \nu = 0$ – Static shell on a Minkowski background

$$\begin{aligned} J_{0+} &= C_{1+} + C_{2+}r^2 + \frac{C_{3+}}{r} + \frac{C_{4+}}{r^3} \\ U_{0+} &= \frac{2\beta_{0+}}{r} + 2C_{2+}r + \frac{2C_{3+}}{r^2} - \frac{3C_{4+}}{r^4} \\ w_{0+} &= -10\beta_{0+}r + 12C_{1+}r - 6C_{2+}r^3 - \frac{6C_{4+}}{r^2} + C_{5+} \end{aligned} \quad (40)$$

where the C_{i+} , $i = 1 \cdots 5$, are constants of integration. An equivalent solution, with constants of integration C_{i-} , $i = 1 \cdots 5$ and with β_{0+} replaced by zero, is applicable in the region $r < r_0$. When we substitute the above solution

into the constraints, we find that $R_{01} = 0$ is satisfied identically, and that, after removing common factors, the other constraints reduce to

$$R_{00} : 4r(2\beta_{0+} - 3C_{1+}) - C_{5+} = 0, \quad q^A R_{0A} : C_{5+} = 0. \quad (41)$$

We require that the solution be regular at null infinity, which means that J must be bounded as $r \rightarrow \infty$, so that

$$C_{2+} = 0. \quad (42)$$

The constraints lead to $C_{1+} = \frac{2}{3}\beta_{0+}$, and since C_{1+} is just $J_{0\infty}$, this leads directly to the news $N = 0$, as expected. In the case that the spacetime includes the origin, the metric must be regular at $r = 0$ so that

$$C_{3-} = C_{4-} = 0. \quad (43)$$

We set the jump in β_0 at $r = r_0$ to be b_c , and we allow arbitrary jumps in w_0 and $J_{0,r}$. The jump constants, together with the remaining unknown $C_{i\pm}$, are fixed by requiring continuity in the metric variables as well as satisfaction of the constraints. We find that the jump conditions at $r = r_0$ are

$$\beta_{0+} = b_c, \quad U_{0+} = U_{0-}, \quad U_{0+,r} = U_{0-,r} + \frac{2b_c}{r_0^2}, \quad w_{0+} = w_{0-} - 2r_0 b_c, \quad J_{0+} = J_{0-}, \quad J_{0+,r} = J_{0-,r}, \quad (44)$$

with the jump in U_0 due to the term $\delta\beta_{,r}$ in the $q^A R_{1A}$ equation, and not sourced by the matter. We have now constructed a metric that is Ricci flat everywhere except on the shell $r = r_0$, and we find the matter source by constructing the Einstein tensor on the shell and using $T^{\alpha\beta} = G^{\alpha\beta}/(8\pi)$. Every component of $T^{\alpha\beta}$ is zero *except*

$$T^{00} = \delta(r - r_0) Z_{\ell m} b_c / (2\pi r_0). \quad (45)$$

If we now set $b_c = 2\pi r_0 \rho_0$, we can interpret $T^{\alpha\beta}$ as being the stress energy tensor of a dust shell with energy density $\rho = \delta(r - r_0) Z_{\ell m} \rho_0$.

B. Case $\nu = 0$, $M \neq 0$ – Static shell on a Schwarzschild background

$$\begin{aligned} J_{0+} &= 2(2M^2 - r^2)C_{1+} + 2 \left(2M^2 - 2rM - 2r^2 + (2M^2 - r^2) \ln \left(1 - \frac{2M}{r} \right) \right) C_{2+} + C_{3+} + \frac{2C_{4+}}{r} + \frac{\beta_{0+}}{2} \\ U_{0+} &= \frac{2\beta_{0+}}{r} - 4(r - 2M)C_{1+} - 4\frac{C_{2+}}{r} \left(2r^2 - 2M^2 - 2rM + r(r - 2M) \ln \left(1 - \frac{2M}{r} \right) \right) + \frac{4C_{4+}(r + M)}{r^3} \\ w_{0+} &= -4\beta_{0+}r + 12C_{1+}r(r - 2M)^2 + 12C_{2+}r \left(2(r^2 + M^2 - 3rM) + (r - 2M)^2 \ln \left(1 - \frac{2M}{r} \right) \right) \\ &\quad + 12C_{3+}r + \frac{12MC_{4+}}{r} + C_{5+} \end{aligned} \quad (46)$$

When we substitute the above solution into the constraints, we find that $R_{01} = 0$ is satisfied identically, and that, after removing common factors, the other constraints reduce to

$$R_{00} : 2r(\beta_{0+} - 6C_{3+}) + 32M^3C_{2+} + 8M\beta_{0+} - C_{5+} = 0, \quad q^A R_{0A} : 32M^3C_{2+} + 8M\beta_{0+} - C_{5+} = 0. \quad (47)$$

The asymptotic form of J_{0+} is

$$J_{0+}(r) \asymp -2(C_{1+} + 2C_{2+})r^2 + 4M^2(C_{1+} + 2C_{2+}) + C_{3+} + \frac{\beta_{0+}}{2}, \quad (48)$$

so that regularity at infinity will require

$$C_{1+} + 2C_{2+} = 0, \quad (49)$$

and thus $J_{0\infty} = C_{3+} + \beta_{0+}/2$. Using the constraints, it follows that $C_{3+} = \beta_{0+}/6$, so that $J_{0\infty} = 2\beta_{0+}/3$, and this immediately leads to the news N being zero as expected.

The jump conditions and stress-energy tensor at $r = r_0$ were determined as in Sec. IV A. The jump conditions are

$$\begin{aligned}\beta_{0+} &= b_c, \quad U_{0+} = U_{0-}, \quad U_{0+,r} = U_{0-,r} + \frac{2b_c}{r_0^2}, \\ w_{0+} &= w_{0-} - 2b_c(r_0 - 2M), \quad J_{0+} = J_{0-}, \quad J_{0+,r} = J_{0-,r} - \frac{Mb_c}{r_0(r_0 - 2M)}.\end{aligned}\quad (50)$$

The only non-zero components of $T^{\alpha\beta}$ are

$$T^{00} = \rho \left(1 - \frac{2M}{r_0}\right)^{-1}, \quad h_{AB}T^{AB} = \rho \frac{M}{r_0^3} \left(1 - \frac{2M}{r_0}\right)^{-1}, \quad q_A q_B T^{AB} = \delta^2 \rho \frac{M}{4r_0^3} \left(1 - \frac{2M}{r_0}\right)^{-1}, \quad (51)$$

where we have set

$$b_c = 2\pi r \rho_0 \left(1 - \frac{2M}{r_0}\right)^{-1} \quad \text{and} \quad \rho = \delta(r - r_0) Z_{20} \rho_0, \quad (52)$$

so that $T^{00} = \rho(v^0)^2$ and thus ρ may be interpreted as an energy density. The above stress-energy tensor is *not* that of a fluid, but represents a rigid shell in which the stresses are such that they preserve the equilibrium of the structure and prevent it from falling towards the black hole. (The above form, in which the density varies with the angle, may be unfamiliar, but the spherically symmetric situation is more commonplace. In this case, ρ in Eq. (52) is replaced by $\rho = \delta(r - r_0)\rho_0$, T^{00} and $h_{AB}T^{AB}$ take the forms given in Eq. (51), and $q_A q_B T^{AB} = 0$; such a stress-energy tensor describes common objects like a balloon, or a bathysphere immersed in the sea.) Returning to the form in Eqs. (51) and (52), we note that the stress-energy tensor has been constructed from a metric, and therefore the Bianchi identities imply $T_{;\beta}^{\alpha\beta} = 0$; we have checked explicitly that this identity is indeed satisfied. Since Z_{20} is negative in places, the stress-energy tensor Eq. (51) violates standard energy conditions [29]. However, that difficulty is easily rectified by supposing that the matter source is a sum of Z_{00} and Z_{20} modes, with the coefficient of Z_{00} larger by a factor of $\sqrt{5}$ than that of Z_{20} . Then, provided $r_0 > 2M$, all the standard energy conditions are satisfied.

There are 10 arbitrary constants available, and these must be chosen so as to satisfy the following 9 conditions

- Regularity at the horizon $r = 2M$ imposes 1 condition, $C_{2-} = 0$;
- Applying the constraints in the exterior imposes the 2 conditions in Eqs. (47) – because the jump conditions preserve the constraints, no additional information is obtained by imposing the constraints in the interior;
- Regularity at infinity imposes 1 condition, Eq. (49);
- There are 5 jump conditions at $r = r_0$.

Thus, we construct a solution that contains 1 arbitrary constant. The values of the various metric quantities at $r = 2M$ are

$$U_{0-} = \frac{3C_0}{2M^2}, \quad U_{0-,r} = -\frac{7C_0}{4M^3} + \frac{\Phi}{M^2}, \quad w_{0-} = 6C_0, \quad J_{0-} = \frac{C_0}{M} + \Phi \quad (53)$$

where C_0 is the arbitrary constant, and where

$$\Phi = \frac{\beta_{0+}}{12M^3} \left((-3r_0^3 + 9r_0^2 M - 12r_0 M^2 + 6M^3) \ln \left(1 - \frac{2M}{r_0}\right) - 14M^3 + 12r_0 M^2 - 6r_0^2 M \right). \quad (54)$$

Thus, we may impose that both of U_{0-} and w_{0-} , or one of $U_{0-,r}$ or J_{0-} , vanish at the horizon, but we *cannot* require

$$U = w = U_{,r} = J = 0 \quad \text{at} \quad r = 2M \quad (55)$$

C. Case $\nu \neq 0$, $M = 0$ – Dynamic shell on a Minkowski background

$$\begin{aligned}J_{0+} &= \frac{C_{1+}}{4r} - \frac{C_{2+}}{12r^3} + \frac{e^{2i\nu r} C_{3+}}{4r^3} (r^2 \nu^2 + 2i\nu r - 1) + \frac{iC_{4+}}{\nu} \\ U_{0+} &= \frac{C_{1+}}{2r^2} + \frac{C_{2+}}{12r^4} (3 + 4i\nu r) + \frac{e^{2i\nu r} C_{3+}}{4r^4} (3 - 2i\nu r) + C_{4+} + \frac{2\beta_{0+}}{r} \\ w_{0+} &= \frac{C_{2+}}{2r^2} (1 + 2i\nu r) + \frac{3e^{2i\nu r} C_{3+}}{2r^2} + 6rC_{4+} \left(\frac{2i}{\nu} - r \right) + C_{5+} - 10\beta_{0+} r\end{aligned}\quad (56)$$

When we substitute the above solution into the constraints, we find that $R_{01} = 0$ is satisfied identically, and that, after removing common factors, the other constraints reduce to

$$R_{00} : 24r\beta_{0+}\nu + 3i\nu^2rC_{1+} - 3\nu^3C_{2+} - 36irC_{4+} + (ir\nu^2 - 3\nu)C_{5+} = 0, \quad q^A R_{0A} : C_{5+} + \nu^2C_{2+} = 0. \quad (57)$$

Regularity at null infinity leads to

$$C_{3+} = 0. \quad (58)$$

Imposing the condition that the spacetime should include the origin means that a series expansion about $r = 0$ of Eqs. (56) must be regular. This leads to

$$C_{1-} = \nu^2C_{3-}, \quad C_{2-} = -3C_{3-}. \quad (59)$$

The jump conditions at $r = r_0$ are

$$\begin{aligned} \beta_{0+} &= b_c, \quad U_{0+} = U_{0-}, \quad U_{0+,r} = U_{0-,r} + \frac{2b_c}{r_0^2} - \frac{4i\nu b_c}{3r_0^2}, \\ w_{0+} &= w_{0-} - 2b_cr_0, \quad J_{0+} = J_{0-}, \quad J_{0+,r} = J_{0-,r} + \frac{r_0\nu^2b_c}{3}. \end{aligned} \quad (60)$$

The only non-zero components of $T^{\alpha\beta}$ are

$$T^{00} = f, \quad q_A T^{0A} = \frac{i\nu\partial f}{6}, \quad q_A q_B T^{AB} = -\frac{\nu^2\partial^2 f}{12}, \quad \text{where } f = \frac{b_c\delta(r-r_0)Z_{20}e^{i\nu u}}{2\pi r_0}. \quad (61)$$

In this case, as expected, the matter shell is dynamic with movement of matter within the shell (because $T^{0A} \neq 0$); but we do not investigate further the physical properties of the matter shell Eq. (61).

D. Case $\nu \neq 0$, $M \neq 0$ – Dynamic shell on a Schwarzschild background

We were not able to obtain a solution in simple analytic form. However, we did obtain series solutions about the singular points of Eq. (39), i.e. about $r = 2M$ and $r = \infty$.

1. Solution about $r = 2M$

Writing $r = y + 2M$, we found that the series solution about $y = 0$ of Eq. (39) contains one singular solution and one regular solution. We discarded the singular solution, and found

$$J_{0-} = -C_{3-}\frac{y^2}{8M^2} \left(1 - \frac{14iM\nu - 3}{3M(4iM\nu - 3)}y + \dots\right) + \frac{C_{2-}}{y + 2M} + C_{1-}. \quad (62)$$

$$U_{0-} = -C_{1-}\nu i + C_{2-}\frac{2(y + 3M)}{(y + 2M)^3} - C_{3-}\frac{1 - 2iM\nu}{2M} \left(1 - \frac{y}{M} + \dots\right) \quad (63)$$

$$w_{0-} = C_{1-}6y(2 + i\nu y + 4iM\nu) - C_{2-}\frac{3y}{y + 2M} + C_{5-} + C_{3-}6(1 - 2i\nu M)y \left(1 + \frac{y^2}{12M^2} + \dots\right) \quad (64)$$

The constraints were evaluated and found to be series with somewhat complicated coefficients, so at first sight it appeared to be impossible to satisfy them. However, the various coefficients are not linearly independent, and impose only two conditions on the constants C_{1-} , C_{2-} , C_{3-} and C_{5-} . Taking C_{1-} and C_{3-} as free, we found that the following solution for C_{5-} and C_{2-} satisfies all the constraints at least up to the order tested (i.e., $\mathcal{O}(y^7)$)

$$C_{5-} = -C_{3-}\frac{3(44iM^2\nu^2 - 8M\nu + 5i + 16M^3\nu^3)}{2\nu(2 + M i\nu)} - C_{1-}\frac{3(8M^3\nu^3 - 30iM^2\nu^2 - 31M\nu + 6i)}{\nu(2 + M i\nu)} \quad (65)$$

$$C_{2-} = -C_{1-}\frac{3i}{\nu} + C_{3-}\frac{-38M\nu - 15i + 24iM^2\nu^2 + 16M^3\nu^3}{6\nu(2 + M i\nu)} \quad (66)$$

2. Solution about $r = \infty$

Here we use $r = 1/x$ and the differential equation to be solved is Eq. (39). This has an irregular singularity at $x = 0$, but nevertheless we were able to construct one regular series solution. Using standard techniques we then constructed the second solution for J_2 , but found, as might be expected, that this solution is singular at $x = 0$; therefore we discarded it. The non-singular solution is

$$J_{0+} = C_{1+} + C_{2+}x + C_{3+}\frac{x^3}{6}\left(1 - \frac{9iM}{4\nu}x^2 + \dots\right). \quad (67)$$

$$U_{0+} = 2\beta_{0+}x - C_{1+}i\nu + C_{2+}2x^2(1 + Mx) - C_{3+}x^3\frac{2i\nu}{3}\left(1 - \frac{3i}{4\nu}x + \dots\right) \quad (68)$$

$$w_{0+} = -10\beta_{0+}x^{-1} + C_{1+}6(2x^{-1} + i\nu x^{-2}) + C_{2+}6Mx - C_{3+}2i\nu x\left(1 - \frac{i}{2\nu}x + \dots\right) + C_{5+}. \quad (69)$$

The situation concerning the constraints is similar to that noted in Sec. IV D 1. Here we take C_{1+} and C_{2+} as free, and find that the following solution satisfies all the constraints up to the order tested (i.e., $\mathcal{O}(x^9)$)

$$C_{5+} = -\beta_{0+}\frac{4(\nu M - 6i)}{\nu} - C_{1+}\frac{18(2i - \nu M)}{\nu} - 12C_{2+} \quad (70)$$

$$C_{3+} = \beta_{0+}\frac{6(2i - \nu M)}{\nu^3} + C_{1+}\frac{9(\nu M - 2i)}{\nu^3} - C_{2+}\frac{3(2 + i\nu M)}{\nu^2} \quad (71)$$

3. Comments

The number of conditions and degrees of freedom is as in the case $\nu = 0$, $M \neq 0$. Thus we expect that it is not possible to impose $J = U = U_{,r} = w = 0$ at $r = 2M$, unless the solution is identically zero throughout the interior; we have confirmed by direct calculation that this is indeed the case.

The coefficients of the series, about both $r = 2M$ and $r = \infty$, have been calculated term by term, and general formulas for the n th coefficient are not known. While the various series do appear to have non-zero radii of convergence, it is not possible to make any formal statement about this issue.

In order to work out the jump conditions and consequently the form of the stress-energy tensor on the shell, we will need to match the series solutions, about $r = 2M$ and $r = \infty$, there. Most likely this will have to be done numerically and is deferred to further work.

V. CONCLUSION

A. Discussion

In Sec. IV B it was shown that, under the circumstances stated, boundary conditions Eq. (55) may not be imposed. Further, Sec. IV D 3 indicates that this result extends to the dynamic, i.e. $\nu \neq 0$, case. Since the imposition of such boundary conditions is common practice [1, 4, 5, 14], the implications of the result need further discussion.

First, let us be clear about the conditions under which the result applies.

1. There needs to be a matter source that is not spherically symmetric around a Schwarzschild black hole. The matter source must be either static, or undergoing regular oscillations ($\propto e^{i\nu u}$).
2. All the metric coefficients need to be either static or undergoing regular oscillations.

Let us note here that the matter source is crucial to the argument, because a vacuum perturbation of Schwarzschild cannot be static or be undergoing regular oscillations: the “no-hair” theorem indicates that such a spacetime must relax to pure Schwarzschild (or, in general, to Kerr if the initial perturbation contains angular momentum).

We investigate further by supposing that condition 1 above applies but not condition 2, and for simplicity we consider the static case. We make the *gedenken* experiment that a spherically symmetric shell has existed around

a black hole since $u = -\infty$. At $u = 0$ the matter starts to move and a new static configuration, with ρ a sum of Z_{00} and Z_{20} modes, is in place from $u = M$ until $u = +\infty$. The situation is illustrated in Fig. 1. For $u < 0$, $\beta = U = U_{,r} = w = J = 0$ at the boundary as well as everywhere inside the shell. For $u > 0$, changes to the gravitational field propagate away from the shell at the speed of light, and the metric relaxes towards a form that satisfies Eq. (53) at $r = 2M$. The inner boundary of the coordinate system is $r = 2M$ on the past event horizon $\mathcal{H}^-$ and is an ingoing null surface. Thus the effects of the shell changing cannot reach it, and the metric $g_{\alpha\beta}$ is well-behaved on each null cone $u = \text{constant}$. However,

$$\lim_{u \rightarrow \infty} g_{\alpha\beta}$$

is not continuous at $r = 2M$. Thus we conclude that if only condition 1 above applies, then homogeneous boundary conditions Eq. (55) may be applied for a finite time lapse u , but not for an infinite or semi-infinite time.

In a numerical simulation using homogeneous boundary conditions Eq. (55) and condition 1, the relaxation of the metric towards the boundary conditions Eq. (53), means that the metric variables must make a finite jump between, say, $r = 2M$ and $r = 2M + \Delta r$. Now, $\Delta r \rightarrow 0$ as $u \rightarrow \infty$, so for short evolutions there is no problem with homogeneous boundary conditions; but for longer evolutions, once Δr becomes smaller than the grid discretization, the solution will seem to be discontinuous and inaccuracies will develop thus making code predictions unreliable.

The difficulty with imposing appropriate boundary data at a black hole horizon in the Bondi-Sachs formalism does not, to our knowledge, affect any results already in the literature. As already stated, the problem arises only if matter is present, and so for example it does not apply to computations involving the scattering of gravitational radiation by a Schwarzschild black hole [14]. Further, the effect comes into play only as $u \rightarrow \infty$, and so does not apply to [4] where the computations terminate at $u = \mathcal{O}(M)$. The earlier work on particle evolution [5] does have long term evolutions, but these were not presented as physical results.

B. Opportunities for further work

As described in Sec. I, the results obtained here are expected to be useful in the validation and improvement of numerical codes using the Bondi-Sachs formalism, particularly in the computation of gravitational radiation. On this point, we note that the coefficient of $J_{0\infty}$ in Eq. (31) scales, for large ℓ , as ℓ^4 , and so it may be that noise at high angular frequency is being multiplied by a large number. Further investigation and numerical testing is required, the starting point for which would probably be the solution exterior to the shell of Sec. IV C ($\nu \neq 0, M = 0$).

This paper has raised the question as to the appropriate boundary data at the horizon for a physical situation involving matter near a Schwarzschild black hole. However, except for the specific case of a static distribution of matter in the form of a thin shell, the issue has not been resolved. Clearly, a prescription is needed when there is a general, dynamic matter distribution.

It is in principle possible to develop the solution obtained in Sec. IV D ($\nu \neq 0, M \neq 0$) to yield the gravitational radiation emitted by a star in close orbit around a black hole, because in this case the metric variables can be expressed as a sum of terms of the form Eq. (19). While this has already been accomplished within a different formalism [30, 31], an alternative derivation would nevertheless be useful. As noted in Sec. IV C, we have not investigated in any detail the physical properties of a time-varying matter shell; it would be necessary to do so in order to pursue this project.

Evaluation of the constraints ($R_{0\alpha} = 0$ in vacuum) has been an essential feature of this paper. Numerical codes based on the characteristic formalism use the constraints in setting boundary data on the inner worldtube – for example homogeneous boundary data (Eq. 55)) trivially satisfy the constraints, and the constraints have been used [6], in a dynamic situation, to determine part of the data. In ADM type codes it is common practice to evaluate the constraints in the interior of the spacetime in order to monitor the accuracy of the evolution. Characteristic codes do not normally evaluate the constraints off the boundary (an exception being [32]), but perhaps this could lead to useful insights.

Acknowledgments

I wish to thank Shrirang Deshingkar, Luis Lehner and Jeffrey Winicour for discussions. I thank Horace Hearne Laboratory of Theoretical Physics, Louisiana State University, as well as Max-Planck-Institut für GravitationsPhysik, Albert-Einstein-Institut, for hospitality. The work was supported in part by the National Research Foundation, South

Africa, under Grant number 2053724, and by the NSF under grant nsf-int 0242507 to Louisiana State University.

-
- [1] N.T. Bishop, R. Gomez, L. Lehner, M. Maharaj and J. Winicour, Phys. Rev. D **56**, 6298 (1997).
 - [2] R. Gómez, Phys. Rev. D **64**, 024007 (2001).
 - [3] N.T. Bishop, R. Gomez, L. Lehner and J. Winicour, Phys. Rev. D **54**, 6153 (1996).
 - [4] N.T. Bishop, R. Gomez, L. Lehner, M. Maharaj and J. Winicour, Phys. Rev. D **60**, 024005 (1999).
 - [5] N.T. Bishop, R. Gomez, S. Husa, L. Lehner and J. Winicour, Phys. Rev. D **68**, 084015 (2003).
 - [6] R. Gomez, S. Husa, L. Lehner and J. Winicour, Phys. Rev. D **66**, 064019 (2002).
 - [7] N.T. Bishop C.J.S. Clarke and R.A. d’Inverno, Class. Quant. Grav. **7**, L23 (1993).
 - [8] R.A. d’Inverno and J.A. Vickers, Phys. Rev. D **56**, 772 (1997).
 - [9] R.A. d’Inverno, M.R. Dubal and E.A. Sarkies, Class. Quant. Grav. **17**, 3157 (2000).
 - [10] P. Papadopoulos and J. A. Font, Phys. Rev. D **61**, 024015 (2000).
 - [11] F. Siebel, J. A. Font, E. Muller and P. Papadopoulos, Phys. Rev. D **67**, 124018 (2003).
 - [12] F. Siebel, J. A. Font and P. Papadopoulos, Phys. Rev. D **65**, 024021 (2002).
 - [13] R. Gomez, L. Lehner, R. L. Marsa and J. Winicour, Phys. Rev. D **57**, 4778 (1998).
 - [14] Y. Zlochower, R. Gomez, S. Husa, L. Lehner and J. Winicour, Phys. Rev. D **68**, 084014 (2003).
 - [15] N.T. Bishop and S.S. Deshingkar, Phys. Rev. D **68**, 024031 (2003).
 - [16] Y. Zlochower, Ph. D. thesis, University of Pittsburgh (2002).
 - [17] J. Winicour, Jnl. Math. Phys. **24**, 1193 (1983).
 - [18] J. Winicour, Jnl. Math. Phys. **24**, 2506 (1983).
 - [19] R.A. Isaacson, J.S. Welling and J. Winicour, Jnl. Math. Phys. **26**, 2859 (1985).
 - [20] M. Campanelli, R Gomez, S. Husa, J. Winicour and Y. Zlochower, Phys. Rev. D **63**, 124013 (2001).
 - [21] S. Husa, Y. Zlochower, R Gomez and J. Winicour, Phys. Rev. D **65**, 084034 (2002).
 - [22] S. A. Teukolsky, Astrophys. J. **185**, 635 (1973).
 - [23] R.A. Isaacson, J.S. Welling and J. Winicour, Jnl. Math. Phys. **24**, 1824 (1983).
 - [24] H. Bondi, M.J.G. van der Burg and A.W.K. Metzner, Proc. R. Soc. **A269**, 21 (1962).
 - [25] R.K. Sachs, Proc. R. Soc. **A270**, 103 (1962).
 - [26] R. Gómez, L. Lehner, P. Papadopoulos and J. Winicour, Class. Quant. Grav. **14**, 977 (1997).
 - [27] E.T. Newman and R. Penrose, Jnl. Math. Phys. **7**, 863 (1966).
 - [28] J.N. Goldberg, A.J. MacFarlane, E.T. Newman, F. Rohrlich and E.C.G. Sudarshan, Jnl. Math. Phys. **8**, 2155 (1967).
 - [29] S.W. Hawking and G.F.R. Ellis, *The large scale structure of spacetime*, (Cambridge University Press, Cambridge, 1974).
 - [30] K. Glampedakis, S. A. Hughes and D. Kennefick, Phys. Rev. D **66**, 064005 (2002).
 - [31] S. A. Hughes, Phys. Rev. D **61**, 084004 (2000).
 - [32] R. Bartnik and A. Norton, <http://www.arXiv.org/gr-qc/9904045> (1999).

FIG. 1: The Penrose diagram of the spacetime described in Sec. V A. $\mathcal{H}^-$ and $\mathcal{H}^+$ are the past and future horizons of the Schwarzschild black hole. The matter shell at $r = r_0$ is the straight line between I^- and I^+ . The line becomes dotted at $u = 0$ indicating a change from spherical symmetry, and the thin dotted line going to $\mathcal{H}^+$ indicates the inner boundary of the domain of influence of this change.